






Population growth of a generational cohort of the copepod *Apocyclops panamensis* (Marsh, 1913) under different temperatures and salinities

Crecimiento poblacional de una cohorte del copépodo *Apocyclops panamensis* (Marsh, 1913) bajo diferentes salinidades y temperaturas.

Leonardo Cruz-Rosado¹ ,
Wilfrido M. Contreras-Sánchez¹ ,
Ulises Hernández-Vidal^{1*} ,
Juan Carlos Perez-Urbiola² ,
María de J Contreras-García¹ 

¹División Académica de Ciencias Biológicas, Universidad Juárez Autónoma de Tabasco, km. 0.5 Carretera Villahermosa-Cárdenas s/n entronque a Bosques de Saloya, CP. 86039, Villahermosa, Tabasco, México.

²Centro de Investigaciones Biológicas del Noroeste, P.O. Box 128, La Paz, B.C.S. 23000, México.

*Corresponding author:
uliseshv44@hotmail.com

Scientific article

Received: march 05, 2020

Accepted: august 13, 2020

How to cite: Cruz-Rosado L, Contreras-Sánchez WM, Hernández-Vidal U, Perez-Urbiola JC, Contreras-García MJ (2020) Population growth of a generational cohort of the copepod *Apocyclops panamensis* (Marsh, 1913) under different temperatures and salinities. *Ecosistemas y Recursos Agropecuarios* 7(2): e2505. DOI: 10.19136/era.a7n2.2505

ABSTRACT. This study was carried out to identify the effects of salinity and temperature during *Apocyclops panamensis* culture, fed with *Tetraselmis chuii*. The experiment was conducted in laboratory conditions for 14 days. Nauplii from the same cohort were used to start the experiment, using a density of 1 ind mL⁻¹ in a volume of 400 mL. The feed consisted of a daily supply of 20 000 cell mL⁻¹ of microalgae. Nine treatments with three replicates were evaluated (24°C-28‰, 24°C-32‰, 24°C-36‰, 28°C-28‰, 28°C-32‰, 28°C-36‰, 32°C-28‰, 32°C-32‰ and 32°C-36‰). The factorial ANOVA indicates that temperature had a significant effect on the abundance of total copepods at day 14 (p = 0.02), while salinity did not show a significant effect (p = 0.06) on the total population of *A. panamensis*. Significant effects of temperature (p < 0.01) and salinity (p < 0.001) were observed for the nauplii population. The best growth of the entire population was reached in the treatment 32°C-28‰, with a population of 1 380.95 (± 1 267.06 ind L⁻¹) at the end of the experiment. Treatments 32°C-28‰ and 32°C-32‰ produced the highest average population of nauplii and copepodites at the end of the experiment. Adult females presented the greatest abundance in the 32°C-32‰ treatment with 214.29 ind L⁻¹, adult males increased between days 11 and 14 in treatments 24°C-28‰, 32°C-28‰, and 32°C-32‰ without exceeding 170 ind L⁻¹. In this study, it is evident that *A. panamensis* prefers warm temperatures and brackish environments.

Key words: Copepod culture, environment requirements, zooplankton.

RESUMEN. El objetivo del estudio fue identificar los efectos de la salinidad y la temperatura durante el cultivo de *Apocyclops panamensis*, alimentado con *Tetraselmis chuii*. El experimento se llevó a cabo en condiciones de laboratorio durante 14 días. Se utilizaron nauplios de una misma cohorte, utilizando una densidad de 1 ind mL⁻¹ en un volumen de 400 mL. La alimentación consistió en un suministro diario de 20 000 células de microalga mL⁻¹. Se evaluaron nueve tratamientos con tres réplicas (24°C-28‰, 24°C-32‰, 24°C-36‰, 28°C-28‰, 28°C-32‰, 28°C-36‰, 32°C-28‰, 32°C-32‰ y 32°C-36‰). El ANOVA factorial indica que la temperatura tuvo un efecto significativo en la abundancia de copéodos totales en el día 14 (p = 0.02), mientras que la salinidad no mostró un efecto significativo (p = 0.06) en la población total de *A. panamensis*. Se observaron efectos significativos de la temperatura (p < 0.01) y la salinidad (p < 0.001) en la población de nauplios. El mejor crecimiento de toda la población al final del experimento se alcanzó en el tratamiento 32°C-28‰, con una población de 1 380.95 ± 1 267.06 ind L⁻¹. Los tratamientos 32°C-28‰ y 32°C-32‰ produjeron la mayor población promedio de nauplios y copepoditos al final del experimento. Las hembras adultas presentaron la mayor abundancia en el tratamiento 32°C-32‰ con 214.29 ind L⁻¹, los machos adultos aumentaron entre los días 11 y 14 en los tratamientos 24°C-28‰, 32°C-28‰, y 32°C-32‰ sin exceder 170 ind L⁻¹. Los resultados indican que *A. panamensis* prefiere las temperaturas cálidas y los ambientes salobres.

Palabras clave: Cultivo de copépodos, requerimientos ambientales, zooplancton.

INTRODUCTION

Planktonic organisms are the basis of the food chain in marine and continental water bodies (Boltovskoy 1981). In marine aquaculture, the use of zooplankton as live food is essential and takes on greater importance during the larval phase, highlighting that it is the best nutritional alternative for the larvae, providing higher survival (Das *et al.* 2012). In the marine environment, copepods are the most dominant, abundant and widely distributed group of microcrustaceans on the planet, sometimes representing up to 80% of the total plankton (Melo *et al.* 2014).

In marine fish culture, rotifers of the genus *Brachionus* are the most used conventional live food for feeding larvae. As they are very active and small-sized organisms, they attract the attention of their predators, allowing them to be easily captured. In addition, by having a high reproduction rate, short life cycle and feeding based on algae, their cultivation is considerably facilitated (Prieto and Atencio 2008, Onyango 2019, Fuller 2020). However, these organisms do not necessarily complete the nutritional requirements for the optimal development of the marine fish larvae that consume them, generating limited growth. All this can lead to high mortality, possibly due to incomplete nutrition and poor development of the digestive system of the larvae (Luna-Figueroa and Arce 2017, Luna-Figueroa *et al.* 2018).

Mass production of copepods has been proposed as an alternative to good quality live food in aquaculture (Suárez-Morales 2000, Prieto *et al.* 2006, Suárez-Morales *et al.* 2009, Rasdi and Qin 2016, Hansen 2017, Martínez-Silva 2018, Hill *et al.* 2020). This proposal takes up the principle that the nutritional value of most copepods is high, and their movement patterns provoke a strong food response in larvae of diverse predatory fish species. Other advantages of using copepods as food are high digestibility, small size, and tolerance to good densities during culture. The simple fact that they are part of the natural food of larvae in the wild allows their use in marine fish culture to promote increases in growth, survival, and quality of the larvae that consume them (Prieto *et al.* 2006,

Martínez-Silva 2018). Nonetheless, when working in aquacultural facilities, it is important to consider that both, rotifers enriched with fatty acids and copepods administered to the fish larvae, increase the possibility of larval survival. The copepods can be considered a nutritional supplement rich in fatty acids.

The quantity, temporality, size, and quality of live food during the larval period are critical to the successful production of marine fish fingerlings. An example of a culture strategy that combines several of these aspects is the so-called mesocosm culture, which has shown promising results by offering a variety of prey that facilitates the larval food transition by imitating what occurs in the wild, and where copepods are a fundamental element (Prieto *et al.* 2006, Rasdi and Qin 2016, Vu *et al.* 2017, Wang *et al.* 2017). The relationship between potential prey and larval selectivity can be approximated by the spatial-temporal coincidence of the two organisms at the spawning sites. An example of this is that the sea herring (*Clupea harengus*), the red snapper (*Lutjanus campechanus*), in its larval stage coexists with a high abundance of zooplankton (Álvarez-Fernández *et al.* 2015, Grüss *et al.* 2018, Heyman *et al.* 2019). The Gulf of Mexico (GM) is no exception because it is also considered an important marine fish spawning area determined by its abundance of larvae (Flores-Coto *et al.* 2009, Heyman *et al.* 2019). Zavala-García *et al.* (2016) recorded the largest zooplankton biomass occurring in GM locations during the same season when spawning grounds of *Centropomus* spp where reported by Hernández-Vidal *et al.* (2014). This event suggests that some of the zooplankton species present, could play an essential role in nutrition and consequently in larval recruitment in the area.

The copepod *Apocyclops panamensis* is a species present in these spawning areas on the coasts of the Gulf of Mexico, so that may conform part of the diet of fish larvae in the region. Preliminary studies indicate that it has the potential to be used as live food since it is resistant to manipulation in captivity, been proposed as a candidate for mass cultivation (Phelps *et al.* 2005, Lindley *et al.* 2011, Santhosh *et al.* 2015, Hill *et al.* 2020). However, the environmental conditions for developing the biotech-

nology for cultivation on a pilot scale are still unknown. For this reason, the objective of this work was to define the effects of salinity and temperature on the population growth of *A. panamensis* under an aquaculture management scheme that could optimize and eventually allow to scale-up production and prove its effectiveness as live food for marine fish species present in the same distribution range.

MATERIALS AND METHODS

This research was carried out in the facilities of the Tropical Aquaculture Laboratory (TAL) of the Biological Sciences Academic Division at the Universidad Juárez Autónoma de Tabasco during August and September of 2019.

Collection and preliminary cultivation of zooplankton

Three zooplankton collection trips were made to the coastal zone in Jalapita, municipality of Centla, Tabasco, Mexico, identified as centropomid spawning areas (Gilmore *et al.* 1983, Hernández-Vidal *et al.* 2014). The collection sites covered an area ranging from 18° 28' 34.71" N - 92° 58' 7.34" W to 18° 26' 26.43" N - 93° 8' 4.58" W, the furthest collection site is 4.7 km from the coastline, and the nearest is approximately 0.2 km from the mouth of the Mecocacán coastal lagoon. Surface trawls (1-2 m deep) were conducted using a 120 μm mesh plankton net. The zooplankton was placed in 500 mL⁻¹ capacity glass jars with water from the sampling site and then moved to the TAL at low temperature in an icebox. In the laboratory, the living material was screened with a 500 μm sieve to remove larger organisms and predators. The remaining organisms were placed in two fiberglass culture tanks with a capacity of 70 L and were maintained at an average temperature of 25.7 \pm 1.13 °C and pH of 7.96 \pm 0.05 using seawater at the same salinity of the collection site (32‰). The water was previously filtered with 2 μm and treated for 10 min with a commercial solution of sodium hypochlorite to reach the concentration of 45ppt and later neutralized by adding 250 mg of sodium thiosulfate L⁻¹. The cultures were fed daily

with a mixture of the microalgae *Nannochloropsis oculata*, *Tetraselmis chuii*, and *T. suecica* supplying a total of 8 million cells per experimental unit, equivalent to a density of 20 000 cells mL⁻¹. After 20 days of culture, the prevailing zooplankton was recovered with a 50 μm sieve that was again maintained for 30 days in conditions similar to those described above.

Isolation and identification

The isolation of the prevailing copepod species began by separating adult specimens. These organisms were concentrated using a 50 μm sieve and under the stereoscopic microscope, and ten adult organisms were isolated. The isolated organisms were kept in 1L-flasks and fed with the previously mentioned algal mixture. This separation of adult organisms was carried out on three occasions, and the organisms were reseeded in new flasks. Samples were fixed in 96% alcohol added with two drops of glycerin in 5mL-tubes. Subsequently, they were sent for determination to the Department of Systematics and Aquatic Ecology at El Colegio de la Frontera Sur (ECOSUR) Chetumal Unit, who determined that the isolated species was *Apocyclops panamensis*. The primary culture was obtained by progressively increasing the volume from the 1L-culture flasks and maintaining the culture conditions previously described.

Experimental design

To evaluate population growth, a completely randomized factorial experiment was carried out with three temperature levels (24, 28, and 32 °C) and three salinity levels (28, 32, and 36‰), resulting in nine treatments, each with six replicates. The experimental units consisted of 0.5L capacity glass flasks containing a working volume of 400 mL. The culture containers were disinfected by washing with a 0.01% sodium hypochlorite solution. To achieve and maintain desired temperatures, the experimental units were placed inside 100L polyethylene tanks, equipped with two 500-watt and controlled with a thermostat (AQUA-KRIL[®] 4160). Seawater (35‰) was used, previously treated with sodium hypochlorite, and neutralized with sodium thiosulfate. The seawater

ter was diluted with sterilized freshwater to reach the salinity of 28 and 32‰, to adjust to 36‰, commercial sea salt was used (Coral Pro-Red Sea[®]). All experimental units were maintained under a photoperiod of 12h:12h (light-dark) controlled with an automatic timer (TEMP 24-HE, STEREN[®]). pH and water temperatures were measured daily using a probe (EcoSense pH10A, YSI USA), while a refractometer was used to measure salinity (Aquafauna[™] Bio-Marine Inc.). Ammonium (NH₃), nitrites (NO₂), and nitrates (NO₃) concentration were measured at the beginning and, subsequently, every five days by colorimetric kits using an aquaculture photometer (HI 83203, HANNA Instruments).

Copepod stocking

A cohort of nauplii was obtained from the main population and separated using a 200 μm mesh sieve to discard copepodites and adult organisms. The nauplii were placed in a glass beaker with sterilized seawater and the density reached was estimated. Before stocking, the seeds were first acclimated to the desired salinity by decreasing or increasing one unit per hour, and then to temperature, by placing the nauplii in 1L flask one hour prior to stocking in each tank with the corresponding temperature. Each experimental unit was stocked with one ind mL⁻¹ for a total of 400 individuals per bottle based on the recommendations of Velásquez *et al.* (2001).

Feeding

The feeding was carried out using the microalgae *Tetraselmis chuii*, according to Velásquez *et al.* (2001). Initially, each unit was supplied with a total of 8 million cells, equivalent to a density of 20 000 cells mL⁻¹. Cultures were used in their maximum growth phase (5-6 days), and the algal density was maintained by counting twice a day and adjusting the preset density when necessary. Microalgae cultures were produced in the Live Food Production Area of the Tropical Aquaculture Laboratory (LAT) at DACBioI using commercial culture medium (PROLINE[®]), the cultures were maintained at constant illumination with fluorescent lamps from 2 000 to 2 500 lux and at a temperature of 24 ± 1.0 °C.

Counting

Counts of copepods were performed every third day, taking seven one mL samples from each experimental unit. The sample was counted using a Bogorov counting chamber (Wildlife Supply model 1810-B20EI) under a stereoscopic microscope (ZEISS[®] model Stemi DV4). The stages identified in the culture count were nauplius, copepodite, non-ovigerous female, ovigerous female, and adult male. In addition, the eggs present in the ovarian sacs of the females were counted. The last evaluation of the population was carried out on the 14th day after sowing, which concluded the experiment.

Statistical analysis of data

The number of organisms per stage was calculated for each treatment, and with the sum of all stages, the value of total copepods (TC) per liter was estimated. After normality and homogeneity of variances were verified (Bartlett's test), the counts obtained at the end of the experiment were compared using a factorial analysis of variance (ANOVA). Significance was determined at the 95% confidence level (α = 0.05). All ANOVAs were performed using the software Statgraphics Centurion XVIII[®]. The graphical representation of the data was done in SigmaPlot v.11[®] package. Results are presented in average values per Liter ± Standard Deviation.

RESULTS

Environmental conditions during the experiment

The environmental conditions recorded throughout the experiment were favorable for the development of copepods (Table 1). The temperature varied slightly around the experimental values. The average temperatures prevailing along the experimental units were 24.24 ± 0.19, 27.86 ± 0.43, and 31.75 ± 0.27. We managed to keep salinity close to the values designated for the study (28 ± 0.18, 32 ± 0.34 and 36 ± 0.32 by monitoring twice a day and replacing evaporated water.

Table 1. Seawater parameters (Mean \pm DE) observed under different temperatures (T) and Salinity (Sal). Nitrites (NO₂), nitrates (NO₃), and total ammonia nitrogen (TAN) were measured every third day, pH, and dissolved oxygen (DO) were measured daily.

T (°C)	Sal (ppm)	NO ₂ (mg L ⁻¹)	NO ₃ (mg L ⁻¹)	TAN (mg L ⁻¹)	pH (IU)	DO (mg L ⁻¹)
24	28	2.30 \pm 1.54	2.47 \pm 4.27	0.37 \pm 0.34	8.12 \pm 0.08	7.17 \pm 0.33
	32	0.67 \pm 0.58	4.87 \pm 4.24	0.49 \pm 0.58	8.11 \pm 0.06	6.60 \pm 0.08
	36	0.33 \pm 0.58	0.63 \pm 1.10	0.47 \pm 0.55	8.10 \pm 0.05	6.54 \pm 0.12
28	28	2.47 \pm 2.54	1.33 \pm 2.31	0.36 \pm 0.35	8.17 \pm 0.09	6.99 \pm 0.24
	32	1.97 \pm 0.95	1.77 \pm 1.91	0.51 \pm 0.50	8.15 \pm 0.05	6.92 \pm 0.30
	36	3.33 \pm 2.08	2.17 \pm 2.20	0.81 \pm 0.95	8.18 \pm 0.08	6.36 \pm 0.16
32	28	1.67 \pm 2.08	1.47 \pm 2.04	0.34 \pm 0.30	8.23 \pm 0.22	5.80 \pm 0.42
	32	1.33 \pm 1.50	1.27 \pm 2.19	0.65 \pm 0.61	8.19 \pm 0.09	6.30 \pm 0.18
	36	2.33 \pm 3.21	3.57 \pm 3.10	0.56 \pm 0.55	8.09 \pm 0.22	6.10 \pm 0.11

Total counts

Results from the factorial ANOVA indicate that the temperature had a significant effect on the overall abundance of organisms in the population assessed at day 14 ($p = 0.02$) and suggests that the evaluated salinities did not have a significant effect on *A. panamensis* production ($p = 0.06$). In this analysis no effect of the interaction of the factors studied was observed ($p > 0.05$). The effect of temperature was that, at a temperature of 32 °C, more organisms were obtained (888.89 ± 414.28 ind L⁻¹) being significantly lower at temperatures of 24 and 28 °C (293.65 ± 285.72 and 357.14 ± 264.70 ind L⁻¹, respectively) (Figure 1A). Although salinity was not statistically significant, a distinctive pattern of decrease in the total number of copepods (TC) can be observed as salinity increases, resulting in $722.22 (\pm 428.57)$, $500.00 (\pm 357.14)$ and $317.46 (\pm 214.29)$ ind L⁻¹ for salinities of 28, 32 and 36‰, respectively (Figure 1B). At the level of specific treatments, the results of TC indicate that the best productivity of this species at the end of the experiment is reached in treatments 32°C-28‰, with a production of $1\,502.00 \pm 457.74$ ind L⁻¹ and 32°C-32‰ where 904.76 ± 576.17 ind L⁻¹ were obtained (Figure 2). When evaluating the effects on the number of copepodites, non-ovigerous females, ovigerous females and males at the end of the experiment, no statistically significant effects of the factors evaluated or their interaction were observed ($p > 0.10$ in all cases). The specific observations for each group of organisms allowed to follow the trends that contribute so that the treatments 32°C-28‰ and 32°C-32‰ have reached the highest total populations at the end

of the experiment.

Nauplii

The factorial ANOVA for the number of nauplii indicated significant effects of temperature ($P = 0.006$) and salinity ($p = 0.001$), with no interaction between these factors ($p > 0.10$). The 32°C-28‰ treatment had the highest average number of nauplii recorded at the end of the experiment, reaching an average of 738.09 ± 602.15 ind L⁻¹. The rebound of this treatment began on day eight, when the population reached 452.38 ind L⁻¹, by day eleven, the population already had $1\,142.86$ ind L⁻¹. This trend was also observed in the 32°C-32‰ treatment, although it was not as pronounced as the previous treatment. In the rest of the treatments, the highest populations did not exceed 400 nauplii L⁻¹, the lowest results being those associated with temperatures of 24°C (Figure 3A).

Copepodites

In general, all treatments showed ups and downs along the trial. The only tendency of increase was observed in treatment 32°C-28‰ for day 11, having an average of 190.47 ± 233.28 ind L⁻¹ and reaching 333.33 ± 147.54 ind L⁻¹ on day 14. All other treatments remained below 150 ind L⁻¹ (Figure 3B).

Adult females

The trend analysis of females includes non-ovigerous females and ovigerous females, showing high variability in the recorded data. The highest




Figure 1. Mean values (\pm Standard error) of total copepods observed in the three temperatures evaluated (A) and the three salinities (B). Different letters indicate statistically significant differences ($p < 0.05$) between temperatures. The sample size per group is $n = 18$.




Figure 2. Treatment trends showing mean values of total copepods observed in the nine treatments evaluated. The sample size per observation is $n = 6$.

values at the end of the experiment correspond once again to treatments 28°C-32‰ and 32°C-28‰ averaging 166.67 ± 167.01 and 119.04 ± 140.46 ind L⁻¹, respectively. The rest of the treatments presented less than 75 ind L⁻¹ (Figure 4A).

Adult males

The number of adult males remained constant

throughout the experiment, with consistent increases observed towards days 11 and 14 in three treatments (24°C-28‰, 32°C-28‰, and 32°C-32‰) however, none of them exceeded 170 ind L⁻¹. The 24°C-28‰ treatment presented the lowest average number of males with $23.81 (\pm 3.04)$ ind L⁻¹ on day 14 of the experiment (Figure 4B).




Figure 3. Treatment trends showing mean values of nauplii (A) and copepodites (B) observed in the nine treatments evaluated for 14 days. The sample size per observation is $n = 6$.




Figure 4. Treatment trends showing mean values for total females (A) and males (B) observed in the nine treatments evaluated for 14 days. The sample size per observation is $n = 6$.

Female:male ratio

This ratio fluctuated considerably throughout the experiment. During the first five days, no adults were observed in the samplings; these were differentiated from day eight. The highest proportion of females was observed in treatments 28 °C-32‰ and 32 °C-28‰ reaching up to six females per male. These values decreased on day 11, with the highest proportion in the treatment being 28 °C-32‰ with five females per male. At the end of the experiment (day 14), the ratio female: male varied in treatments

between 0.29 and 3.0, averaging between treatments 1.1 females per male.

DISCUSSION

The population trends observed during 14 days starting from a cohort of nauplii indicate that *Apocyclops panamensis* grow well under temperatures near 32°C and salinities ranging from 28 and 32‰. *A. panamensis* is one of the few species of cy-

clopoids cultured in the laboratory and recommended for aquacultural practices; however, despite this recommendation, little is known about the specific environmental requirements for this copepod species abundant in the neotropical region (Suárez-Morales *et al.* 2004). The fact that the two best treatments correspond to the temperature of 32°C indicates that this species has better performance under the upper range of temperatures measured in warm-tropical environments. In recent years, we have recorded an average of 29.5°C for the warmest months of the year (September-October) near the sites where *A. panamensis* was collected for the current study, and this corresponds a shallow coastal habitat where is possible to reach high temperature during sunny and wind calm days. Our study suggests that *A. panamensis* like most copepods display a wide tolerance range for temperature fluctuations.

There is a variety of information regarding the effects of temperature on copepod populations, but the leading theory focuses on the direct effects of temperature on fecundity and growth. Warm temperatures promote a fast life cycle, which can translate into high population growth in a short period; this is possible because the duration of the developmental stages and molting rates are physiological processes that are primarily dictated by temperature (Hirst and Bunker 2003, Chailinn *et al.* 2018). Proof of this is that many tropical Cyclopoid copepods have life cycles lasting from four to five days, making them susceptible to mass cultivation (Su *et al.* 2005, Rasdi *et al.* 2018). While some species can acclimatize to temperatures outside their normal ranges without inhibiting their reproductive potential (Verbitsky *et al.* 2017), other species make reproductive adjustments, for example, *Acartia tonsa* - a temperate climate species - decreases the number of eggs produced by females in cold temperatures (Hansen *et al.* 2010). Similar results have been reported for *A. distans*, a species belonging to the same genus, whose highest population growth was achieved under a temperature range of 26 to 33.8°C.

Salinity is another of the environmental parameters that strongly influence the ecological and biological responses of copepods (Pan *et al.* 2016).

In this experiment, it becomes evident that *A. panamensis* is well adapted to a relatively wide range of salinities; however, population growth displays best results at 28‰, particularly for the naupliar stage. This species is considered euryhaline since it has been reported in environments ranging from the oligohaline condition that occurs in coastal lagoons and estuaries to hypersaline waters in semi-desert areas of North America (Pérez *et al.* 2006, Reid and Hribar 2006, Annabi-Trabelsi *et al.* 2019). The wide tolerance of the species has also been demonstrated by being collected in salinity of 6.6‰ and later successfully acclimatized to 30‰ (Lindley *et al.* 2011, Roy and Venkataraman 2018). It is inferred that it is a species that tolerates wide ranges of salinity, with a preference for brackish water as observed in this experiment. Even though euryhaline species can survive under a wide range of salinities, reproductive performance can be affected severely depending on this parameter leading to unsustainable population growth and ultimately result in copepod mortality (Pan *et al.* 2016). Reid *et al.* (2002) and Fuentes-Reinés and Suárez-Morales (2015) considered that the genus *Apocyclops* is so ubiquitous that can be observed dominating plankton communities in coastal lagoons, saline lakes, and coastal marshes. However, it is frequently associated with salinities under 30‰. In our experiment, all treatments having 32‰ resulted in reduced production of nauplii. This response to higher salinity levels is explained by Kimmel and Bradley (2001) and Pan *et al.* (2016) as a result of a shift in the energy budget for reproduction towards energy needed for adjusting the metabolism required for osmoregulation. The fact that we have registered population growth in the various conditions evaluated is proof of the vast adaptive capacity of *A. panamensis*.

In general for this group of organisms, it is known that adults have tolerance to several factors considered adverse as well as their latency stages since this facilitates dissemination by means of birds and wind, which could be the explanation of the successful colonization of contrasting environments in which they have been found, although little is known about the physiological and ecological mechanisms

that allow copepods to reside in these environments (Anufriieva 2015).

In the best treatment of this study (32°C-28‰), it was observed that the population grew 3.4 times in 14 days, reflecting rapid growth. These results are similar to other studies where Cyclopoid production has commonly been generated from 1 to 5 times the amount inoculated (Pérez *et al.* 2006, Farhadian *et al.* 2008, Ruiz-Guzmán *et al.* 2012). In an extreme case, Velásquez *et al.* (2001) reported a growth in the culture of *A. distans* of up to 27 times what was stocked for ten days using ambient temperature (26.0 - 38.8°C), salinity around 40‰ and total darkness for their best treatment. The high productivity observed in our treatments mainly reflects the number of nauplii obtained, which are the objective for the aquaculture of fish larvae.

The obtained productivity of 738 nauplii L⁻¹ is a moderate nauplii production when compared to values obtained by Phelps *et al.* (2005) in the experimental culture of *A. panamensis* using adult copepods in initial densities of 320, 1 280 and up to 5 120 adults L⁻¹ for a period of 4 to 9 days. In that study, the author produced 16 942 nauplii L⁻¹ in the maximum density of adults stocked. These differences could be attributable to culture procedures and stages stocked, but suggest potential for nauplii production in *A. panamensis* under our defined environmental conditions.

Another factor that could have contributed to the productivity observed in *A. panamensis* is the type of food used. *Tetraselmis chuii* is considered an excellent food alternative for copepods, because they are mobile flagellated algae of considerable size, having a notorious preference in comparison with non-mobile algae. Velásquez *et al.* (2001) obtained the best production values of nauplii in *A. distans* using *Tetraselmis chuii* when compared with *N. oculata*. The cell density used in this study can be considered as low (20 000 cell mL⁻¹) compared to other studies where 50,000 cell mL⁻¹ (Pérez *et al.* 2006) to 600 000 cell mL⁻¹ (Mujica *et al.* 1995) are used. Velásquez *et al.* (2001) used algae densities of 300 000 cell mL⁻¹ of *T. chuii* to feed *A. distans*, while Ruiz-Guzmán *et al.* (2012) obtained the best results

in the cultivation of *Cyclopina* sp. with the combination of *T. suecica* and *Isochrysis galbana*. The cell density of 600 000 cell mL⁻¹, gave favorable results in the culture of *Tigriopus* sp. supplying *Nannochloropsis* sp. (Mujica *et al.* 1995). Phelps *et al.* (2005) cultivated *A. panamensis* using a cell density of 500 000 cell mL⁻¹ using the microalgae *Isochrysis galbana*. The production of *A. panamensis* could be improved by using a varied diet, possibly favoring the productivity of copepod crops compared to feeding using a single species. This can be attributed to the high content of polyunsaturated fatty acids provided by the mixtures (Farhadian 2008). Alternatively, these algal diets can be improved by supplying other food sources such as flours of vegetable and animal origin, as well as inorganic fertilizers and using culture protocols such as mesocosm resulting in a good nutritional profile (Phelps *et al.* 2005, Prieto *et al.* 2006).

Copepods play an important role in the initial stages of development of marine fish because they constitute their main natural prey, in addition to having a high nutritive content, easy digestion and adequate transfer of nutrients, making them an important food source in aquaculture (Ruiz-Guzmán *et al.* 2012). Copepod nauplii are a rich source of free amino acids (FAA) and contain more than twice the amount of FAA than *Artemia*, and higher levels of HUFAs (Naess *et al.* 1995, Rayner *et al.* 2017). Proof of this nutritional value is that copepods are used in the larval stage of turbot *Scophthalmus maximus* (Bruno *et al.* 2018) Atlantic cod *Gadus morhua* (Karlsen *et al.* 2015), red snapper *Lutjanus campechanus* (Phelps *et al.* 2005) and the gilt-head sea bream *Sparus aurata* (Mona *et al.* 2019). All these species proved to be successful predators of all stages of copepods of the orders Harpacticoida and Cyclopoida, demonstrating the potential of copepod cultivation by improving the nutritional quality of the diet and decreasing dependence on rotifers and artemia (Vanacor-Barroso *et al.* 2017). Data for FAA content in *A. panamensis* indicate that nutritional profile is similar to other copepod species used actually on larviculture (Lindley *et al.* 2011).

Copepods can be cultivated at high densities, and that their cultivation is relatively easy (Ribeiro and Souza-Santos 2011). They have even been

successfully cultivated in recirculation systems where the production has been exclusively of nauplii and copepodites, stages that are used in the feeding of marine fish larvae (Buttino *et al.* 2012). In a recent study, we found that *A. panamensis* maintained at 28‰ and 32°C accomplishes its life cycle in six days (data not shown) and reached 11 600 copepods L⁻¹ in a 1 000 L tank. With these preliminary results, *A. panamensis* can be considered as a good candidate for mass culture protocol evaluations in order to determine conditions for pilot-scale production (Ribeiro and Souza-Santos 2011, Hansen *et al.* 2017). Previous experiences by Phelps (2005) in *A. panamensis* cultured in earthen ponds indicate feasibility for mass production on this species; however, more information about the optimal conditions of variables in the culture system is necessary. The results presented here provide valuable information for *A. panamensis* culture, because the definition of preferred temperatures and salinities for a particular species, provide conditions where the organisms are subject to minimum stress; their physiological functions are optimized, providing a maximum population growth (Nichelmann 1983, Verbitsky *et al.* 2017). These data can be suitable for production protocols evaluations.

CONCLUSIONS

The species *Apocyclops panamensis* has a preference for warm waters (32°C) and medium salinity (28-32‰). The high production capacity of

A. panamensis observed in this experiment allows visualizing the potential for scaling up production and evaluation as live food for larval culture of tropical fish. Nauplii and copepodites can be an alternative for feeding larval stages of marine fishes due to their small sizes. Another convenient feature is that it has a short life cycle that makes them susceptible to rapid mass production. We consider that *A. panamensis* is a species suitable to be used as live food; however, it is required to demonstrate its viability for mass production at pilot scale and preference by larvae of different species of fish.

ACKNOWLEDGEMENTS

Authors thank the anglers from the Sociedad Cooperativa de Pescadores “San Ramón” Tabasco, especially Ramón Domínguez-Sánchez for his invaluable logistic support during our biological samplings. This investigation was a part of the Aquaculture Collaborative Research Support Program, supported partly by the United States Agency for Internal Development (USAID) (LAG-G-00-96-90015-00). The opinions expressed herein are those of the authors and do not necessarily reflect the views of the AquaFish Innovation Lab or USAID. CONA-CyT provided a Ph.D. scholarship to Leonardo Cruz-Rosado (258602). W.M.C.S. is a grant holder from SNI-CONACyT.

LITERATURE CITED

- Álvarez-Fernández S, Licandro P, Van Damme CJG, Hufnagl M (2015) Effect of zooplankton on fish larval abundance and distribution: a long-term study on North Sea herring (*Clupea harengus*). *ICES Journal of Marine Science* 72: 2569-2577.
- Annabi-Trabelsi N, El-Shabrawy G, Goher ME, Subrahmanyam MNV, Al-Enezi Y, Ali M, Ayadi H, Belmonte G (2019) Key Drivers for Copepod Assemblages in a Eutrophic Coastal Brackish Lake. *Water* 11: 2-20.
- Anufrieva EV (2015) Do copepods inhabit hypersaline waters worldwide? A short review and discussion. *Chinese Journal of Oceanology and Limnology* 33: 1354-1361.
- Boltovskoy D (1981) Atlas del Atlántico Sudoccidental y métodos de trabajo con el zooplancton marino. Mar del Plata. INIDEP. Argentina. 936p.

- Bruno E, Højgaard JK, Hansen BW, Munk P, Støttrup JG (2018) Influence of swimming behavior of copepod nauplii on feeding of larval turbot (*Scophthalmus maximus*). *Aquaculture International* 26: 225-236.
- Buttino I, Ianora A, Buono S, Vitiello V, Malzone MG, Rico C, Langellotti AL, Sansone G, Gennari L, Miralto A (2012) Experimental cultivation of the Mediterranean calanoid copepods *Temora stylifera* and *Centropages typicus* in a pilot re-circulating system. *Aquaculture Research* 43: 247-259.
- Chailinn P, Ju SJ, Park W, Kim HW, Lee SR, Park JH (2018) The strategy of population maintenance by coastal copepod inferred from seasonal variations in abundance of adults and resting eggs. *Ocean and Polar Research* 40: 213-222.
- Das P, Mandal SC, Bhagabati SK, Akhtar MS, Singh S K (2012) Important live food organisms and their role in aquaculture. *Frontiers in Aquaculture* 4: 69-86.
- Farhadian O, Yusoff FM, Arshad A (2008) Population growth and production of *Apocyclops dengizicus* (Copepoda: Cyclopoida) fed on different diets. *Journal of World Aquaculture Society* 39: 384-396.
- Flores-Coto C, Espinosa-Fuentes M, Zavala-García F, Sanvicente-Añorve L (2009) Ictioplancton del sur del Golfo de México: Un compendio. *Hidrobiológica* 19: 49-76
- Fuentes-Reinés JM, Suárez-Morales E (2015) Checklist of planktonic copepoda from a colombian coastal lagoon with record of *Halicyclops exiguus* kiefer. *Boletín de Investigaciones Marinas y Costeras - INVEMAR* 44: 369-389.
- Fuller SA (2020) Minimizing time fed rotifers maximizes hybrid striped bass larval growth in recirculating aquaculture systems. *North American Journal of Aquaculture* 2: 208-214.
- Gilmore RG, Donohoe CJ, Cooke DW (1983) Observations on the distribution and biology of east-central Florida populations of the common snook, *Centropomus undecimalis* (Bloch). *Florida Scientist* 46: 313-336.
- Grüss A, Biggs C, Heyman WD, Erisman B (2018) Prioritizing monitoring and conservation efforts for fish spawning aggregations in the US Gulf of Mexico. *Fisheries Research* 213: 75-93.
- Hansen B W, Drillet G, Kozmér A, Madsen KV, Pedersen MF, Sørensen TF (2010) Temperature effects on copepod egg hatching: does acclimatization matter? *Journal of Plankton Research* 32: 305-315.
- Hansen BW (2017) Advances using copepods in aquaculture. *Journal of Plankton Research* 39: 972-974.
- Hansen BW, Hansen PJ, Nielsen TG, Jepsen PM (2017) Effects of elevated pH on marine copepods in mass cultivation systems: Practical implications. *Journal of Plankton Research* 39: 984-993.
- Hernández-Vidal U, Chiappa-Carrara X, Contreras-Sánchez W (2014) Reproductive variability of the common snook, *Centropomus undecimalis*, in environments of contrasting salinities interconnected by the Grijalva-Usumacinta fluvial system. *Ciencias Marinas* 40: 173-185.
- Heyman WD, Grüss A, Biggs CR, Kobara SI, Farmer NA, Karnauskas M, Lowerre-Barbieri S, Erisman B (2019) Cooperative monitoring, assessment, and management of fish spawning aggregations and associated fisheries in the U.S. Gulf of Mexico. *Marine Policy* 109: 103689. Doi: 10.1016/j.marpol.2019.103689.
- Hill M, Pernetta A, Crooks N (2020) Size matters: A review of live feeds used in the culture of marine ornamental fish. *Asian Fisheries Science* 33: 161-174.
- Hirst AG, Bunker AJ (2003) Growth of marine planktonic copepods: global rates and patterns in relation to chlorophyll a, temperature, and body weight. *Limnology Oceanography* 48: 1988-2010.

- Karlsen Ø, Van Der MT, Rønnestad I, Mangor-Jensen A, Galloway TF, Kjørsvik E, Hamre K (2015) Copepods enhance nutritional status, growth and development in Atlantic cod (*Gadus morhua* L.) larvae-can we identify the underlying factors? *PeerJ* 3: e902. Doi. 10.7717/peerj.902.
- Kimmel DG, Bradley BP (2001) Specific protein responses in the calanoid copepod *Eurytemora affinis* (Pope, 1880) to salinity and temperature variation. *Journal of Experimental Marine Biology and Ecology* 266: 135-149.
- Lindley LC, Phelps RP, Davis DA, Cummins KA (2011) Salinity acclimation and free amino acid enrichment of copepod nauplii for first-feeding of larval marine fish. *Aquaculture* 318: 402-406.
- Luna-Figueroa J, Arce UE (2017) Un menú diverso y nutritivo en la dieta de peces: “el alimento vivo”. *Agro Productividad* 10: 112-116
- Luna-Figueroa J, Arce UE, Figueroa TJ (2018) Ventajas e inconvenientes del uso de alimento vivo en la nutrición de peces. *Inventio* 14: 39-43.
- Martínez-Silva MA, Audet C, Winkler G, Tremblay R (2018) Prey quality impact on the feeding behavior and lipid composition of winter flounder (*Pseudopleuronectes americanus*) larvae. *Aquaculture and Fisheries*. 3: 145-155.
- Melo PAMC, de Melo J MDM, Macêdo SJ, Araujo M, Neumann-Leitao S (2014) Copepod distribution and production in a Mid-Atlantic Ridge archipelago. *Anais da Academia Brasileira de Ciências* 4: 1719-1733.
- Mona MH, Rizk EST, El-feky MM, Elawany ME (2019) Effect of nutritional quality of rotifers and copepods on sea bream (*Sparus aurata*) fry fish productivity. *The Egyptian Journal of Experimental Biology* 15: 135-142.
- Mujica AR, Carvajal UC, Miranda EO (1995) Cultivo experimental de *Tigriopus* sp. (Copepoda: Harpacticoidea). *Investigaciones Marinas* 23: 75-82.
- Naess T, Germain-Henry M, Naas KE (1995) First feeding of Atlantic halibut (*Hippoglossus hippoglossus*) using different combinations of *Artemia* and wild zooplankton. *Aquaculture* 130: 235-250.
- Nichelmann M (1983) Some characteristics of the biological optimum temperature. *Journal of Thermal Biology* 8: 69-71.
- Onyango OC (2019) The mass culture of the freshwater rotifers *Brachionus rubens* Ehrenberg 1838 using different Algal species diets *United International Journal for Research & Technology* 1:1-15.
- Pan YJ, Souissi A, Souissi S, Hwang JS (2016) Effects of salinity on the reproductive performance of *Apocyclops royi* (Copepoda, Cyclopoida). *Journal of experimental Marine Biology and Ecology* 475: 108-113.
- Pérez IE, Gómez NS, Puello-Cruz AC (2006) Efecto de diferentes dietas de microalgas sobre la supervivencia y crecimiento de *Apocyclops aff. panamensis* (Marsh, 1913), (Copépodo: Cyclopoida) cultivado bajo condiciones controladas de laboratorio. *Revista Electronica de Veterinaria* 7:1-10.
- Phelps RP, Sumiarsa S, Lipman EE, Lan HP, Moss KK, Davis AD (2005) Intensive and extensive production techniques to provide copepod nauplii for feeding larval red snapper *Lutjanus campechanus*. In: Cheng-Sheng L, O'Bryen P, Marcus N (ed) *Copepods in aquaculture* Wiley-Blackwell. US. pp: 151-168.
- Prieto M, Atencio V (2008) Zooplancton en la larvicultura de peces neotropicales. *Revista MVZ Cordoba* 13:1415-1415.
- Prieto M, Castaño F, Sierra J, Logato P, Botero J (2006) Alimento vivo en la larvicultura de peces marinos: copépodos y mesocosmos. *Revista MVZ Córdoba* 11: 30-36.

- Rasdi NW, Qin JG (2016) Improvement of copepod nutritional quality as live food for aquaculture: a review. *Aquaculture Research* 47: 1-20.
- Rasdi NW, Suhaimi H, Yuslan A, Sung YY, Ikhwanuddin, Omar SS, Qin JG, Kassim Z, Yusoff FM (2018) Effect of mono and binary diets on growth and reproduction of cyclopoid copepod. *Aquaculture, Aquarium, Conservation and Legislation* 11: 1658-1671.
- Rayner TA, Jørgensen NOG, Drillet G, Hansen, BW (2017) Changes in free amino acid content during naupliar development of the calanoid copepod *Acartia tonsa*. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 210: 1-6.
- Reid JW, Hamilton IV R, Duffield RM (2002) First confirmed New World record of *Apocyclops dengizicus* (Lepeleshkin), with a key to the species of *Apocyclops* in North America and the Caribbean region (Crustacea: Copepoda: Cyclopidae). *Jeffersoniana* 10: 1-25.
- Reid JW, Hribar LJ (2006) Records of some Copepoda (Crustacea) from the Florida Keys. *Proceedings of the Academy of Natural Sciences of Philadelphia. Jeffersoniana* 155: 1-8.
- Ribeiro AC, Souza-Santos LP (2011) Mass culture and offspring production of marine harpacticoid copepod *Tisbe biminiensis*. *Aquaculture* 321: 280-288.
- Roy MD, Venkataraman K (2018) Catalogue on copepod fauna of India. part 2. cyclopoida and harpacticoida (arthropoda: crustacea). *Journal of Environment and Sociobiology* 15: 109-194.
- Ruiz-Guzmán JA, Jiménez CA, Gomes C, Prieto MJ (2012) Cultivo experimental de *Cyclopina* sp. con diferentes especies de microalgas. *Revista Colombiana de Ciencias Pecuarias* 25: 97-105.
- Santhosh BF, Anzeer FM, Unnikrishnan C, Anil MK (2015) Potential species of copepods for marine finfish hatchery. In: Vizhinjam Research Centre of CMFRI (ed). *Teaching Resource. Vizhinjam*. pp: 170-176.
- Su HM, Cheng SH, Chen TI, Su MS (2005) Culture of copepods and applications to marine finfish larval rearing in Taiwan. In: Cheng-Sheng L, O'Bryen P, Marcus N (ed) *Copepods in aquaculture*. Wiley-Blackwell US. Pp: 183-194.
- Suárez-Morales E (2000) Copépodos, seres ubicuos y poco conocidos. *Biodiversitas* 29: 7-11.
- Suárez-Morales E, Fleeger JW, Montagna PA (2009) Free-living Copepoda (Crustacea) of the Gulf of Mexico. In: Felder DL, Camp DK, Cato JC, Buster NA, Holmes ChW, Day JW, Yáñez AA, Bianchi TS (ed.) *Gulf of Mexico origin, Waters and Biota*. Vol 1. College Station, Texas, United States : Texas A&M University Press. USA. pp: 841-870.
- Suárez-Morales E, Reid JW, Fiers F, Iliffe TM (2004) Historical biogeography and distribution of the freshwater cyclopine copepods (Copepoda, Cyclopoida, Cyclopinae) of the Yucatan Peninsula, Mexico. *Journal Biogeography* 31: 1051-1063.
- Vanacor-Barroso M, Carvalho CVAD, Antoniassi R, Ronzani-Cerqueira V (2017) The copepod *Acartia tonsa* as live feed for fat snook (*Centropomus parallelus*) larvae from notochord flexion to advanced metamorphosis. *Latin American Journal of Aquatic Research* 45: 159-166.
- Velásquez A, Rosas J, Cabrera T, Millán J, Hernández M (2001) Efecto de *Tetraselmis chuii*, *Nannochloris oculata* y *Dunaliella salina* sobre el crecimiento poblacional de *Apocyclops distans* (Copepoda, Cyclopoidae) en diferentes condiciones de temperatura e iluminación. *Revista de Biología Marina y Oceanografía* 36: 189-197.

- Verbitsky VB, Grishanin AK, Malysheva OA, Medyantseva EN, Verbitskaya TI (2017) Thermal resistance, preferred and avoidance temperatures of *Cyclops strenuus* Fischer, 1851, and their relation to optimal, pessimal, and tolerant temperatures. *Biology Bulletin* 44: 439-448.
- Vu MTT, Hansen BW, Kiørboe T (2017) The constrains of high density production of the calanoid copepod *Acartia tonsa* Dana. *Journal of Plankton Research* 39: 1028-1039.
- Wang KKL, Shao J, HU W (2017) Yeast and corn flour supplement to enhance large-scale culture efficiency of marine copepod *Tisbe furcata*, a potential live food for fish larvae. *Israeli Journal of Aquaculture-Bamidgeh* 69: 1-8.
- Zavala-García F, Flores-Coto C, De la Luz Espinosa-Fuentes M (2016) Relationship between zooplankton biomass and continental water discharges in the southern Gulf of Mexico (1984-2001). *Revista de Biología Marina y Oceanografía* 51: 21-31.